

# Medical Standing Order for NASAL NALOXONE 4 mg

## 2024 - 2025 School Year

# **Opioid Overdose** - Always an **EMERGENCY**

- Naloxone (Narcan) is indicated for reversal of suspected opioid overdose resulting in respiratory depression and/or unresponsiveness to stimuli.
- Person with symptoms of overdose must ALWAYS be transported by EMS to hospital for additional medical attention.

#### Signs of Overdose

- <u>Neuro</u> Unresponsive to stimuli such as shaking or sternal rub, limp body.
  *Pinpoint Pupils*
- Lungs slow, shallow breathing or no breathing.
- <u>Heart</u> Slow or no pulse. Cyanotic coloration to lips and fingertips.
- Mouth unable to speak/incoherent. Gurgling or deep snorting.
- <u>Skin</u> Pale, Clammy. Cyanotic coloration to lips and fingertips.
- <u>GI</u> Vomiting or gurgling.
- <u>Other</u> Report from bystanders, prior history or presence of drug paraphernalia

### Action:

- Call 911
- Institute School Emergency Plan
- Administer NARCAN Nasal Spray
  - > Lay person on back
  - > Tilt head back slightly
  - Insert Narcan nozzle tip gently into ONE NOSTRIL
  - Press Plunger Firm to administer dose
  - Place person in Recovery position

1 | Page 4/18/24 | f



Non-Public School Health Program

- Repeat Dose in 3 5 minutes to other nostril if person does not respond by waking up to voice or touch.
- Monitor vital signs and airway until EMS arrive; if needed start Rescue breathing and CPR
- Individual is to be transported by EMS even if symptoms improve.
- Continue to monitor vital signs and record. If a student, provide a copy of student's Emergency Form / permission to treat.
- If the parents/guardian are not present at time of transport to the hospital, a school representative must accompany the student to the hospital.

Neil Stein, un

4/30/2024

Neil Stein, MD

**Medical Director** 

Date